

Infections in hemodialysis: a concise review. Part II: blood transmitted viral infections.

Eleftheriadis T, Liakopoulos V, Leivaditis K, Antoniadis G, Stefanidis I

Nephrology Department, Medical School, University of Thessaly, Larissa, Greece

Abstract

Hemodialysis (HD) patients are particularly predisposed to infections. It seems that the HD procedure per se as well as disturbances in both innate and adaptive immunity significantly contribute to this susceptibility. Infections are the major cause of morbidity and the second cause of death following cardiovascular events in HD patients. Episodes of bacteremia and pneumonia account for the majority of severe infections in this population. In addition to these bacterial infections another common problem in HD units is the blood transmitted viral infections, particularly infections caused by hepatitis B virus, hepatitis C virus and Human immunodeficiency virus. A number of safety concerns exist for limiting the spread of these viral infections among HD patients and the staff of the unit. The aim of the present review is to present in a concise albeit practical form the difficult aspect of infections in HD. For practical reasons the review is separated in two parts. The previous first part covered bacteremia and respiratory infections, while the present second part covers blood transmitted viral infections. Hippokratia 2011; 15 (2): 120-126

Key words: hemodialysis, infection, hepatitis B virus, hepatitis C virus, human immunodeficiency virus, review

Corresponding author: Theodoros Eleftheriadis, Department of Nephrology, Medical School, University of Thessaly, Neo Ktirio, Mezourlo Hill, 41110 Larissa, Greece, e-mail: teleftheriadis@yahoo.com, Tel: 306977522168, Fax:30 2413500241

Hemodialysis (HD) procedure per se as well as disturbances in both innate¹⁻³ and adaptive immunity⁴⁻⁶ make HD patients susceptible to infections. Infections are the major cause of morbidity and the second cause of death following cardiovascular events in HD patients. Interestingly, death risk from cardiovascular events significantly increases after hospitalization due to infection. Episodes of bacteremia account for the majority of severe infections in this population, while episodes of pneumonia follow⁷. The annual mortality due to bacteremia is 100-300 times higher in HD patients compared to the general population. Even when age, race, sex, diabetes and record errors are taken into account, mortality owing to bacteremia is still 50 times higher^{8,9}. Besides bacterial infections, another common problem in HD units is the blood transmitted viral infections, particularly infections caused by hepatitis B virus (HBV), hepatitis C virus (HCV) and Human immunodeficiency virus (HIV). Due to the nature of the HD procedure, safety concerns exist for limiting their spread among HD patients and the staff of the unit. In addition, the natural history of all these infections, the available treatments and the response to vaccines differ from what is known for the general population. There are many appreciable reviews that analyze rather extensively each infectious agent separately¹⁰⁻¹⁵. The aim of the present review is to present in a concise albeit practical form a global update of the difficult aspect of infections in HD. In the present second part of the review blood transmitted infections, which are present in all HD units, are discussed.

Hepatitis B and hemodialysis

It is estimated that approximately 350 million people are chronic hepatitis B virus (HBV) carriers worldwide¹⁶. Consequently, most HD units treat chronic HBV carriers. Interestingly, because of the known acquired immunity disturbances in this population⁵, after the initial HBV infection, 60% of hemodialysis patients become chronic carriers, while the respective percentage in the general population is only 5%¹⁷⁻¹⁹. Fortunately, HBV carriage does not significantly affect prognosis of HD patients. Although 30% of HBV carriers develop histologically confirmed chronic hepatitis, only 5% die from liver disease^{20,21}. However, life threatening exacerbations and increased rates of liver disease were reported in renal transplant recipients who were asymptomatic during HD²²⁻²⁶. In a meta-analysis chronic HBV infection was associated with an increased risk of death (RR 2.49) and graft loss (RR 1.44)²⁷.

Serology helps to determine the HBV status of HD patients (Table 1). A HBsAg, anti-HBc and anti-HBs negative patient is susceptible and needs vaccination. A HBsAg negative but anti-HBc and anti-HBs positive patient is immune due to natural infection, while a HBsAg and anti-HBc negative but anti-HBs positive patient is immune due to vaccination. HBsAg, anti-HBc and IgM anti-HBc positivity without anti-HBs indicate an acutely infected patient, while HBsAg, anti-HBc positivity without IgM anti-HBc and anti-HBs indicate the chronic carrier state. The situation is more complex when a patient

Table 1: Hepatitis B serology

| Serological markers | Interpretation |
|---|---|
| HBsAg (-) anti-HBc (-) anti-HBs (-) | Susceptibility |
| HBsAg (-) anti-HBc (-) anti-HBs (+) | Immunity because of vaccination |
| HBsAg (-) anti-HBc (+) anti-HBs (+) | Immunity because of natural infection |
| HBsAg (+) anti-HBc (+) anti-HBs (-) IgM anti-HBc (+) | Acute infection |
| HBsAg (+) anti-HBc (+) anti-HBs (-) IgM anti-HBc (-) | Chronic infection |
| HBsAg (-) anti-HBc (+) anti-HBs (-) | False positive anti-HBc test or Resolved acute infection or Resolved infection or "Low level" chronic infection |
| HBeAg (+) HBV DNA (+) | Active virus replication |

is HBsAg and anti-HBs negative but anti-HBc positive. The above serology could mean resolved infection which is the most common, or resolving acute infection, or false positive anti-HBc, or "low level" chronic infection²⁸. The last case is of particular interest and a Canadian study showed that although in 241 hemodialysis patients only 2 (0.8%) were HBsAg positive, among the rest 239 patients, 9 (3.8%) were found to be HBV-DNA positive with PCR²⁹. Thus HBV-DNA test is useful for anti-HBc positive patients who are also negative for HBsAg and anti-HBs. Additionally, nowadays hidden hepatitis is revealed with nested PCR for the pre-S/S, pre-core/core and X regions of the HBV genome³⁰.

When HBV carriage is confirmed, the follow up of HD patients includes serum alanine transaminase (ALT) and aspartate transaminase (AST) every 1-2 months, alpha-fetoprotein (AFP) every 3-4 months, albumin and prothrombin time. The hypoaminotransferasemia that characterize HD patients should be taken into consideration, since the normal ALT and AST levels are less than 27 IU/L and 17 IU/L in this population³¹. Interestingly, the absence of HBeAg does not necessarily exclude viral replication if there is a mutation of the pre-core/core gene inducer³². HBV-DNA should be tested at least annually and liver biopsy is recommended when specific treatment is to be started or prior to renal transplantation^{33,34}.

Significant viral replication, with more than 100000 copies/ml with PCR, indicates the need for treatment³⁵. Despite the limited data, interferon A seems to be beneficial³⁶. However, the increased rate of side-effects in HD patients, mainly anemia and cachexia, establishes nucle-

oside analogues as the treatment of choice^{37,38}. Nucleosides analogues should be administered for at least one year. Lamivudine at a dose of 10mg per day is particularly effective in HD patients, since it suppresses viral replication and transaminasemia in 80% of patients³⁹⁻⁴¹. Unfortunately, 15 % of patients develop resistance after one year of treatment and 30% after two years due to mutation of RNA-dependent HBV-DNA polymerase⁴²⁻⁴⁴. Telvivudine at a dose of 600mg every 96 hours is characterized by much less resistance development compared to lamivudine and can be used as first line treatment⁴⁵. New but not yet tested in HD patients nucleosides analogues remain to be tried.

For preventing intra-unit HBV spread, the usual precautions for blood transmitted infections should be strictly applied. Blood could be present even on surfaces that seem very clean with naked eye⁴⁶. Despite some doubt^{47,48}, patients should be hemodialyzed in separate rooms⁴⁹, since HBV is found in high titers and is potentially infectious for more than 7 days in the environment. In comparison, hepatitis C virus (HCV) is less infectious and survives for a shorter period of time, while Human immunodeficiency virus (HIV) is far less infectious and cannot survive in natural environment⁵⁰⁻⁵². The risk of infection after contaminated needle stick exposure is about 6% if the patient is HBeAg negative and about 30% if the patient is HBeAg positive⁵³. If not vaccinated, which is unacceptable for the HD unit staff, the individual should receive a dose of HBV immune globulin (HBIG) (0.6mg/kg) and start vaccination. If the individual is vaccinated and the anti-HBs levels are higher than 10 IU/ml no measures are required. For the vaccinated individual with anti-HBs levels below 10 IU/ml a dose of HBIG plus a repetitive vaccine dose are indicated. Finally, two HBIG doses are required for individuals who did not respond after two complete series of vaccination⁵⁴.

Vaccination against HBV is also an effective measure, since vaccinated patients have 70% less possibility to become HBV carriers⁵⁵. Unfortunately, due to the impaired immune function of HD patients 3-6⁵⁶, response rates determined as anti-HBs titres higher than 10 IU/L are smaller in HD patients than in the general population. Despite the administration of double vaccine doses and of an extra dose in HD patients, response rate is about 50-60%, while the response rate in the general population is higher than 90%^{57,58}. Some authors referred a higher, but still less than in the general population, vaccine-induced seroconversion rate (75-80%), possibly due to differences in the patients' selection criteria^{56,59}. All doses of the vaccine should be repeated in patients who have not responded 1-2 months after complete first vaccination series⁶⁰. Nevertheless, only one dose is required in patients with initial anti-HBs levels greater than 10 IU/L due to successful response to vaccination or natural infection who now present with reduced anti-HBs levels^{61,62}. Vaccination should be performed as soon as possible in the course of chronic renal failure since response is associated with the degree of renal function⁶³. Finally, new

vaccines against HBV containing new immune adjuvants are promising. For example, the HBV-AS04 vaccine was found to provide better initial response rates (78% vs. 51%) and also greater duration of response during the 42 months study period⁶⁴.

Hepatitis C and hemodialysis

There are about 170 million hepatitis C virus (HCV) carriers worldwide and HD patients belong to a high-risk population⁶⁵. A meta-analysis revealed that in HD patients HCV carriage is associated with 1.57 times increased risk of death. Liver cirrhosis and hepatoma contribute to the increased mortality⁶⁶.

Typically HCV carriage is indicated by the presence of anti-HCV antibodies, which currently are assessed in the serum with third generation ELISA. Optimally, a positive result could be confirmed with RIBA (recombinant immunoblot assay) due to the higher specificity of this assay^{67,68}. Detection of HCV-RNA with PCR also confirms the diagnosis of HCV carriage and indicates viral replication as well. However, it should be noted that in the course of HCV carriage there are intervals without detectable HCV-RNA in the serum⁶⁹. Recently, HCV core Ag detection with ELISA can be used for the diagnosis of HCV carriage⁷⁰⁻⁷².

Interestingly, a patient can be anti-HCV positive, but HCV-RNA negative. Such results could mean that the patient has been cured or reflect interrupted viremia that is common in the course of HCV carriage. A false positive ELISA or a viremia below the PCR lower detection limit could also be responsible⁷³⁻⁷⁵. In the opposite situation, a patient can be anti-HCV negative, but HCV-RNA positive. This could be the result of inability to develop a satisfactory antibody titre due to the immune dysfunction that accompanies HD or simply reflect the window between infection and antibody production. Certainly, a false positive ELISA may also be the cause⁷⁶⁻⁷⁸.

Currently, it is suggested to use HCV-RNA detection for diagnosis of HCV carriage in areas or HD units with high HCV prevalence, while anti-HCV antibodies detection is better to be used in areas or HD units with low HCV prevalence. A positive anti-HCV test should always be followed by a HCV-RNA test. The latter should also be applied in case of hypertransaminasemia even if the anti-HCV test is negative. The screening with the one or the other method should be repeated every 6-12 months⁷⁹.

Antiviral treatment is indicated in patients with chronic kidney disease –besides transplanted patients – with acute HCV infection who did not have negative PCR after 12 weeks. Patients from transplantation lists that meet the above criteria should be treated. Regarding the HCV-infected kidney transplant recipients, it is suggested that treatment should be considered only when the benefits of treatment clearly outweigh the risk of allograft rejection due to interferon A based therapy. In the latter patients, such a treatment could be applied in case of fibrosing cholestatic hepatitis or life-threatening vasculitis. Generally, the desired serologic response to treatment is a nega-

tive HCV-RNA PCR 6 months after treatment is completed. If a desired serologic response is achieved, retesting for HCV-RNA every 6 months is recommended⁸⁰.

Currently, routine ribavirin treatment is contraindicated in HD patients⁸¹ and small trials showed increased rate of adverse effects with pegylated interferon A^{82,83}. Thus the only available treatment is interferon A. A meta-analysis of 20 studies showed that treatment with interferon A led to a desired serological response in 41% of patients, while 26% of them discontinued the treatment because of side-effects. The results were better with doses greater or equal to 3000000 units 3 times per week. However, higher doses were associated with more frequent discontinuation of the drug due to adverse effects. Lower HCV-RNA titers and transaminase levels and absence of cirrhosis at the initiation of treatment were associated with a better prognosis⁸⁴.

In order to prevent the intra-unit spread of HCV, adhesion to general precautions for blood transmitted infections is required. Despite the existing doubts^{85,86}, HCV positive patient isolation or use of separate machines are not officially recommended^{49,87-89}. The risk of infection due to contaminated needle stick exposure is about 1.8 % and unfortunately there is no preventive post exposure treatment available⁵³.

Human immunodeficiency virus and hemodialysis

The prevalence of Human immunodeficiency virus (HIV) positivity varies and is dependent on the region where the dialysis unit is located^{90,91}. In the year 2000 1% of patients on dialysis in the USA has reached end-stage renal disease (ESRD) due to HIV associated nephropathy (HIVAN)⁹². HIVAN shows a predilection for black race^{93,94} and accounts for the majority (65%) but not all of the ESRD cases among HIV positive patients⁹⁵.

All HD patients should be tested for HIV since prompt diagnosis, monitoring and timely initiation of Highly Active Anti-Retroviral Therapy (HAART) consisted of three or more highly potent anti-HIV drugs, commonly reverse transcriptase inhibitors and protease inhibitors, improves prognosis. A positive ELISA must be confirmed with western blotting as in general population. Among patients with positive ELISA and moderately positive western blotting, up to 4.5% of results can be false positive. This possibility is increased in cases of blood transfusions or renal transplantation. It is clear that, in case of positive results, serum HIV-RNA detection is mandatory⁹⁶.

Adhesion to general precautions for blood transmitted infections is adequate for preventing intra-unit spread of HIV and CDC does not recommend patient isolation or separate machines⁴⁹. However if precautions are overlooked the results can be disastrous^{97, 98}. The risk of infection after contaminated needle stick exposure is about 0.3%^{99,100}. Preventive post-exposure treatment should start immediately. It includes 2-3 anti-retroviral drugs for a 4 weeks period and it reduces the risk of infection by 80%. The individual should be checked for HIV 0, 1, 2, 3

and 6 months after exposure¹⁰¹⁻¹⁰³.

In HIV positive HD patients native arteriovenous fistula creation should be encouraged since grafts show increased thrombosis and infection rates in these patients, possibly because of intravenous drug abuse^{104,105}. However, arteriovenous grafts are preferred to CVC due to high infection rates of CVC¹⁰⁶. Anemia is another common problem in these patients. Frequent infections and HAART toxicity have been incriminated¹⁰⁷. Increased cardiovascular risk is another adverse effect of long lasting HAART administration¹⁰⁸.

The prognosis of HIV positive HD patients has significantly improved. HAART¹⁰⁹, disease stage at initiation of dialysis¹¹⁰ and CD4 lymphocyte count¹¹¹ are the major prognostic factors. In 1980 mortality was 100% for the first year¹¹⁰. In the USA, the one-year survival has increased from 56% for those who started dialysis in 1990 to 74% for those who started in 1999¹¹². Finally, in a recent study from France, no difference in one year survival was found between HIV positive and HIV negative patients¹¹³. All of the above benefits are mainly attributed to HAART, which must include the proper drugs given at the right time and at the right doses. Otherwise, control of viremia might be impaired or toxicity can be significant resulting in increased mortality. Cooperation with other specialists is mandatory¹¹⁴⁻¹¹⁶.

References

- Lewis SL, Van Epps DE, Chenoweth DE. Alterations in chemotactic factor-induced responses of neutrophils and monocytes from chronic dialysis patients. *Clin Nephrol.* 1988; 30: 63-72.
- Muniz-Junqueira MI, Braga Lopes C, Magalhaes CA, Schleicher CC, Veiga JP. Acute and chronic influence of hemodialysis according to the membrane used on phagocytic function of neutrophils and monocytes and pro-inflammatory cytokines production in chronic renal failure patients. *Life Sci.* 2005; 77: 3141-3155.
- Eleftheriadis T, Kartsios C, Yiannaki E, Kazila P, Antoniadis G, Liakopoulos V, et al. Chronic inflammation and CD16+ natural killer cell zeta-chain downregulation in hemodialysis patients. *Blood Purif.* 2008; 26: 317-321.
- Eleftheriadis T, Papazisis K, Kortsaris A, Vayonas G, Voytzi S, Vargemzis V. Impaired T cell proliferation and zeta chain phosphorylation after stimulation with staphylococcal enterotoxin-B in hemodialysis patients. *Nephron Clin Pract.* 2004; 96: c15-20.
- Eleftheriadis T, Antoniadis G, Liakopoulos V, Kartsios C, Stefanidis I. Disturbances of acquired immunity in hemodialysis patients. *Semin Dial.* 2007; 20: 440-451.
- Eleftheriadis T, Kartsios C, Yiannaki E, Kazila P, Antoniadis G, Liakopoulos V, et al. Chronic inflammation and T cell zeta-chain downregulation in hemodialysis patients. *Am J Nephrol.* 2008; 28: 152-157.
- Morbidity & Mortality. *American journal of kidney diseases: the official journal of the National Kidney Foundation*; 55: S269-S280.
- Sarnak MJ, Jaber BL. Mortality caused by sepsis in patients with end-stage renal disease compared with the general population. *Kidney Int.* 2000; 58: 1758-1764.
- Foley RN, Guo H, Snyder JJ, Gilbertson DT, Collins AJ. Sepsis in the United States dialysis population, 1991 to 1999. *J Am Soc Nephrol.* 2004; 15: 1038-1045.
- Beathard GA, Urbanes A. Infection associated with tunneled hemodialysis catheters. *Semin Dial.* 2008; 21: 528-538.
- Kausz A, Pahari D. The value of vaccination in chronic kidney disease. *Semin Dial.* 2004; 17: 9-11.
- Hussein MM, Mooij JM, Roujouleh H. Tuberculosis and chronic renal disease. *Semin Dial.* 2003; 16: 38-44.
- Fabrizi F, Messa P, Martin P. Hepatitis B virus infection and the dialysis patient. *Semin Dial.* 2008; 21: 440-446.
- Fabrizi F, Lunghi G, Ganeshan SV, Martin P, Messa P. Hepatitis C virus infection and the dialysis patient. *Semin Dial.* 2007; 20: 416-422.
- Rao TK. Human immunodeficiency virus infection in end-stage renal disease patients. *Semin Dial.* 2003; 16: 233-244.
- Lavanchy D. Hepatitis B virus epidemiology, disease burden, treatment, and current and emerging prevention and control measures. *J Viral Hepat.* 2004; 11: 97-107.
- Szmuness W, Prince AM, Grady GF, Mann MK, Levine RW, Friedman EA, et al. Hepatitis B infection. A point-prevalence study in 15 US hemodialysis centers. *JAMA.* 1974; 227: 901-906.
- London WT, Drew JS, Lustbader ED, Werner BG, Blumberg BS. Host responses to hepatitis B infection in patients in a chronic hemodialysis unit. *Kidney Int.* 1977; 12: 51-58.
- Sampliner RE. The duration of hepatitis B surface antigenemia. *Arch Intern Med.* 1979; 139: 145-146.
- Harnett JD, Parfrey PS, Kennedy M, Zeldis JB, Steinman TI, Guttman RD. The long-term outcome of hepatitis B infection in hemodialysis patients. *Am J Kidney Dis.* 1988; 11: 210-213.
- Josselson J, Kyser BA, Weir MR, Sadler JH. Hepatitis B surface antigenemia in a chronic hemodialysis program: lack of influence on morbidity and mortality. *Am J Kidney Dis.* 1987; 9: 456-461.
- Friedlaender MM, Kasper RT, Rubinger D, Silver J, Popovtzer MM. Renal transplantation is not contraindicated in asymptomatic carriers of hepatitis B surface antigen. *Am J Kidney Dis.* 1989; 14: 204-210.
- Pirson Y, Alexandre GP, Ypersele C. Long-term effect of hbs antigenemia on patient survival after renal transplantation. *N Engl J Med.* 1977; 296: 194-196.
- Weir MR, Kirkman RL, Strom TB, Tilney NL. Liver disease in recipients of long-functioning renal allografts. *Kidney Int.* 1985; 28: 839-844.
- Parfrey PS, Forbes RD, Hutchinson TA, Kenick S, Farge D, Dauphinee WD, et al. The impact of renal transplantation on the course of hepatitis B liver disease. *Transplantation.* 1985; 39: 610-615.
- Chan TM, Fang GX, Tang CS, Cheng IK, Lai KN, Ho SK. Preemptive lamivudine therapy based on HBV DNA level in HBsAg-positive kidney allograft recipients. *Hepatology.* 2002; 36: 1246-1252.
- Fabrizi F, Martin P, Dixit V, Kanwal F, Dulai G. HBsAg seropositive status and survival after renal transplantation: meta-analysis of observational studies. *Am J Transplant.* 2005; 5: 2913-2921.
- Mast EE, Margolis HS, Fiore AE, Brink EW, Goldstein ST, Wang SA, et al. A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP) part 1: immunization of infants, children, and adolescents. *MMWR Recomm Rep.* 2005; 54: 1-31.
- Minuk GY, Sun DF, Greenberg R, Zhang M, Hawkins K, Uhanova J, et al. Occult hepatitis B virus infection in a North American adult hemodialysis patient population. *Hepatology.* 2004; 40: 1072-1077.
- Hui CK, Sun J, Au WY, Lie AK, Yueng YH, Zhang HY, et al. Occult hepatitis B virus infection in hematopoietic stem cell donors in a hepatitis B virus endemic area. *J Hepatol.* 2005; 42: 813-819.
- Yasuda K, Okuda K, Endo N, Ishiwatari Y, Ikeda R, Hayashi H, et al. Hypoaminotransferasemia in patients undergoing long-term hemodialysis: clinical and biochemical appraisal. *Gastro-*

- enterology. 1995; 109: 1295-1300.
32. Blum HE. Hepatitis B virus: significance of naturally occurring mutants. *Intervirology*. 1993; 35: 40-50.
 33. Ozdogan M, Ozgur O, Gur G, Boyacioglu S, Ozderin Y, Demirhan B, et al. Histopathological impacts of hepatitis virus infection in hemodialysis patients: should liver biopsy be performed before renal transplantation? *Artif Organs*. 1997; 21: 355-358.
 34. Degott C, Degos F, Jungers P, Naret C, Courouce AM, Potet F, et al. Relationship between liver histopathological changes and HBsAg in 111 patients treated by long-term hemodialysis. *Liver*. 1983; 3: 377-384.
 35. Gane E, Pilmore H. Management of chronic viral hepatitis before and after renal transplantation. *Transplantation*. 2002; 74: 427-437.
 36. Duarte R, Huraib S, Said R, Abdel-Khadir A, Sullivan S, Chaballout A, et al. Interferon-alpha facilitates renal transplantation in hemodialysis patients with chronic viral hepatitis. *Am J Kidney Dis*. 1995; 25: 40-45.
 37. Chan TM, Wu PC, Lau JY, Lok AS, Lai CL, Cheng IK. Interferon treatment for hepatitis C virus infection in patients on haemodialysis. *Nephrol Dial Transplant*. 1997; 12: 1414-1419.
 38. Campistol JM, Esforzado N, Martinez J, Rosello L, Veciana L, Modol J, et al. Efficacy and tolerance of interferon-alpha(2b) in the treatment of chronic hepatitis C virus infection in haemodialysis patients. Pre- and post-renal transplantation assessment. *Nephrol Dial Transplant*. 1999; 14: 2704-2709.
 39. Boyacioglu S, Gur G, Gursoy M, Ozdemir N. Lamivudine in renal transplant candidates with chronic hepatitis B infection. *Transplant Proc*. 2002; 34: 2131-2132.
 40. Schmilovitz-Weiss H, Melzer E, Tur-Kaspa R, Ben-Ari Z. Excellent outcome of Lamivudine treatment in patients with chronic renal failure and hepatitis B virus infection. *J Clin Gastroenterol*. 2003; 37: 64-67.
 41. Lapinski TW, Flisiak R, Jaroszewicz J, Michalewicz M, Kowalczyk O. Efficiency and safety of lamivudine therapy in patients with chronic HBV infection, dialysis or after kidney transplantation. *World J Gastroenterol*. 2005; 11: 400-402.
 42. Kletzmayr J, Watschinger B, Muller C, Demetriou D, Puchhammer-Stockl E, Ferenci P, et al. Twelve months of lamivudine treatment for chronic hepatitis B virus infection in renal transplant recipients. *Transplantation*. 2000; 70: 1404-1407.
 43. Liaw YF, Leung NW, Chang TT, Guan R, Tai DI, Ng KY, et al. Effects of extended lamivudine therapy in Asian patients with chronic hepatitis B. *Asia Hepatitis Lamivudine Study Group*. *Gastroenterology*. 2000; 119: 172-180.
 44. Fontaine H, Thiers V, Chretien Y, Zylberberg H, Poupon RE, Brechot C, et al. HBV genotypic resistance to lamivudine in kidney recipients and hemodialyzed patients. *Transplantation*. 2000; 69: 2090-2094.
 45. Lai CL, Gane E, Liaw YF, Hsu CW, Thongsawat S, Wang Y, et al. Telbivudine versus lamivudine in patients with chronic hepatitis B. *N Engl J Med*. 2007; 357: 2576-2588.
 46. Bergervoet PW, van Riessen N, Sebens FW, van der Zwet WC. Application of the forensic Luminol for blood in infection control. *J Hosp Infect*. 2008; 68: 329-333.
 47. Tokars JI, Alter MJ, Miller E, Moyer LA, Favero MS. National surveillance of dialysis associated diseases in the United States-1994. *ASAIO J*. 1997; 43: 108-119.
 48. Tokars JI, Finelli L, Alter MJ, Arduino MJ. National surveillance of dialysis-associated diseases in the United States, 2001. *Semin Dial*. 2004; 17: 310-319.
 49. Recommendations for preventing transmission of infections among chronic hemodialysis patients. *MMWR Recomm Rep*. 2001; 50: 1-43.
 50. Bond WW, Favero MS, Petersen NJ, Gravelle CR, Ebert JW, Maynard JE. Survival of hepatitis B virus after drying and storage for one week. *Lancet*. 1981; 1: 550-551.
 51. Favero MS, Bond WW, Petersen NJ, Berquist KR, Maynard JE. Detection methods for study of the stability of hepatitis B antigen on surfaces. *J Infect Dis*. 1974; 129: 210-212.
 52. Kamili S, Krawczynski K, McCaustland K, Li X, Alter MJ. Infectivity of hepatitis C virus in plasma after drying and storing at room temperature. *Infect Control Hosp Epidemiol*. 2007; 28: 519-524.
 53. Beltrami EM, Williams IT, Shapiro CN, Chamberland ME. Risk and management of blood-borne infections in health care workers. *Clin Microbiol Rev*. 2000; 13: 385-407.
 54. Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HBV, HCV, and HIV and Recommendations for Postexposure Prophylaxis. *MMWR Recomm Rep*. 2001; 50: 1-52.
 55. Miller ER, Alter MJ, Tokars JI. Protective effect of hepatitis B vaccine in chronic hemodialysis patients. *Am J Kidney Dis*. 1999; 33: 356-360.
 56. Eleftheriadis T, Liakopoulos V, Antoniadis G, Stefanidis I, Galaktidou G. Indoleamine 2,3-dioxygenase is increased in hemodialysis patients and affects immune response to hepatitis B vaccination. *Vaccine*. 2011; 29: 2242-2247.
 57. Stevens CE, Alter HJ, Taylor PE, Zang EA, Harley EJ, Szmuness W. Hepatitis B vaccine in patients receiving hemodialysis. Immunogenicity and efficacy. *N Engl J Med*. 1984; 311: 496-501.
 58. Buti M, Viladomiu L, Jardi R, Olmos A, Rodriguez JA, Bartolome J, et al. Long-term immunogenicity and efficacy of hepatitis B vaccine in hemodialysis patients. *Am J Nephrol*. 1992; 12: 144-147.
 59. Bruguera M, Rodicio JL, Alcazar JM, Oliver A, Del Rio G, Esteban-Mur R. Effects of different dose levels and vaccination schedules on immune response to a recombinant DNA hepatitis B vaccine in haemodialysis patients. *Vaccine* 1990; 8 Suppl: S47-9; discussion S60-62.
 60. Dinitis-Pensy M, Forrest GN, Cross AS, Hise MK. The use of vaccines in adult patients with renal disease. *Am J Kidney Dis*. 2005; 46: 997-1011.
 61. Charest AF, Grand'Maison A, McDougall J, Goldstein MB. Evolution of naturally acquired hepatitis B immunity in the long-term hemodialysis population. *Am J Kidney Dis*. 2003; 42: 1193-1199.
 62. Tsouchnikas I, Dounousi E, Xanthopoulou K, Papakonstantinou S, Thomoglou V, Tsakiris D. Loss of hepatitis B immunity in hemodialysis patients acquired either naturally or after vaccination. *Clin Nephrol*. 2007; 68: 228-234.
 63. DaRoza G, Loewen A, Djurdjev O, Love J, Kempston C, Burnett S, et al. Stage of chronic kidney disease predicts seroconversion after hepatitis B immunization: earlier is better. *Am J Kidney Dis*. 2003; 42: 1184-1192.
 64. Kong NC, Beran J, Kee SA, Miguel JL, Sanchez C, Bayas JM, et al. A new adjuvant improves the immune response to hepatitis B vaccine in hemodialysis patients. *Kidney Int*. 2008; 73: 856-862.
 65. Yen T, Keeffe EB, Ahmed A. The epidemiology of hepatitis C virus infection. *J Clin Gastroenterol*. 2003; 36: 47-53.
 66. Fabrizi F, Martin P, Dixit V, Bunnapradist S, Dulai G. Meta-analysis: Effect of hepatitis C virus infection on mortality in dialysis. *Aliment Pharmacol Ther*. 2004; 20: 1271-1277.
 67. Fabrizi F, Lunghi G, Raffaele L, Guarnori I, Bacchini G, Corti M, et al. Serologic survey for control of hepatitis C in haemodialysis patients: third-generation assays and analysis of costs. *Nephrol Dial Transplant*. 1997; 12: 298-303.
 68. Fehr T, Ambuhl PM. Chronic hepatitis virus infections in patients on renal replacement therapy. *Nephrol Dial Transplant*. 2004; 19: 1049-1053.
 69. Fabrizi F, Poordad FF, Martin P. Diagnostic workup of hepatitis C and the patient on maintenance dialysis. *Int J Artif Organs*.

- 2001; 24: 843-852.
70. Tanaka E, Ohue C, Aoyagi K, Yamaguchi K, Yagi S, Kiyosawa K, et al. Evaluation of a new enzyme immunoassay for hepatitis C virus (HCV) core antigen with clinical sensitivity approximating that of genomic amplification of HCV RNA. *Hepatology*. 2000; 32: 388-393.
 71. Aoyagi K, Ohue C, Iida K, Kimura T, Tanaka E, Kiyosawa K, et al. Development of a simple and highly sensitive enzyme immunoassay for hepatitis C virus core antigen. *J Clin Microbiol*. 1999; 37: 1802-1808.
 72. Kurtz JB, Boxall E, Qusir N, Shirley J, Coleman D, Chandler C. The diagnostic significance of an assay for 'total' hepatitis C core antigen. *J Virol Methods*. 2001; 96: 127-132.
 73. Galan F, Perez-Gracia MT, Lozano A, Benavides B, Fernandez-Ruiz E, Rodriguez-Iglesias MA. A 3-year follow-up of HCV-RNA viraemia in haemodialysis patients. *Nephrol Dial Transplant*. 1998; 13: 1211-1214.
 74. Barril G, Castillo I, Arenas MD, Espinosa M, Garcia-Valdecasas J, Garcia-Fernandez N, et al. Occult hepatitis C virus infection among hemodialysis patients. *J Am Soc Nephrol*. 2008; 19: 2288-2292.
 75. Umlauf F, Gruenewald K, Weiss G, Kessler H, Urbanek M, Haun M, et al. Patterns of hepatitis C viremia in patients receiving hemodialysis. *Am J Gastroenterol*. 1997; 92: 73-78.
 76. al Meshari K, al Ahdal M, Alfurayh O, Ali A, De Vol E, Kessie G. New insights into hepatitis C virus infection of hemodialysis patients: the implications. *Am J Kidney Dis*. 1995; 25: 572-578.
 77. Kalantar-Zadeh K, Miller LG, Daar ES. Diagnostic discordance for hepatitis C virus infection in hemodialysis patients. *Am J Kidney Dis*. 2005; 46: 290-300.
 78. Pereira BJ, Milford EL, Kirkman RL, Quan S, Sayre KR, Johnson PJ, et al. Prevalence of hepatitis C virus RNA in organ donors positive for hepatitis C antibody and in the recipients of their organs. *N Engl J Med*. 1992; 327: 910-915.
 79. Guideline 1: Detection and evaluation of HCV in CKD. *Kidney Int*. 2008; 73: S10-S19.
 80. Guideline 2: Treatment of HCV infection in patients with CKD. *Kidney Int*. 2008; 73: S20-S45.
 81. Meyers CM, Seeff LB, Stehman-Breen CO, Hoofnagle JH. Hepatitis C and renal disease: an update. *Am J Kidney Dis*. 2003; 42: 631-657.
 82. Teta D, Luscher BL, Gonvers JJ, Francioli P, Phan O, Burnier M. Pegylated interferon for the treatment of hepatitis C virus in haemodialysis patients. *Nephrol Dial Transplant*. 2005; 20: 991-993.
 83. Russo MW, Ghalib R, Sigal S, Joshi V. Randomized trial of pegylated interferon alpha-2b monotherapy in haemodialysis patients with chronic hepatitis C. *Nephrol Dial Transplant*. 2006; 21: 437-443.
 84. Gordon CE, Uhlrig K, Lau J, Schmid CH, Levey AS, Wong JB. Interferon treatment in hemodialysis patients with chronic hepatitis C virus infection: a systematic review of the literature and meta-analysis of treatment efficacy and harms. *Am J Kidney Dis*. 2008; 51: 263-277.
 85. dos Santos JP, Loureiro A, Cendoroglo Neto M, Pereira BJ. Impact of dialysis room and reuse strategies on the incidence of hepatitis C virus infection in haemodialysis units. *Nephrol Dial Transplant*. 1996; 11: 2017-2022.
 86. Jadoul M, Cornu C, van Ypersele de Strihou C. Incidence and risk factors for hepatitis C seroconversion in hemodialysis: a prospective study. The UCL Collaborative Group. *Kidney Int*. 1993; 44: 1322-1326.
 87. Jadoul M, Cornu C, van Ypersele de Strihou C. Universal precautions prevent hepatitis C virus transmission: a 54 month follow-up of the Belgian Multicenter Study. The Universitaires Cliniques St-Luc (UCL) Collaborative Group. *Kidney Int*. 1998; 53: 1022-1025.
 88. Aucella F, Vigilante M, Valente GL, Stallone C. Systematic monitor disinfection is effective in limiting HCV spread in hemodialysis. *Blood Purif*. 2000; 18: 110-114.
 89. Guideline 3: Preventing HCV transmission in hemodialysis units. *Kidney Int*. 2008; 73: S46-S52.
 90. Perez G, Ortiz-Interian C, Lee H, de Medina M, Cerney M, Alain JP, et al. Human immunodeficiency virus and human T-cell leukemia virus type I in patients undergoing maintenance hemodialysis in Miami. *Am J Kidney Dis*. 1989; 14: 39-43.
 91. Vigneau C, Guiard-Schmid JB, Tourret J, Flahault A, Rozenbaum W, Pialoux G, et al. The clinical characteristics of HIV-infected patients receiving dialysis in France between 1997 and 2002. *Kidney Int*. 2005; 67: 1509-1514.
 92. Eggers PW, Kimmel PL. Is there an epidemic of HIV infection in the US ESRD program? *J Am Soc Nephrol*. 2004; 15: 2477-2485.
 93. Choi AI, Rodriguez RA, Bacchetti P, Bertenthal D, Volberding PA, O'Hare AM. Racial differences in end-stage renal disease rates in HIV infection versus diabetes. *J Am Soc Nephrol*. 2007; 18: 2968-2974.
 94. Lucas GM, Lau B, Atta MG, Fine DM, Keruly J, Moore RD. Chronic kidney disease incidence, and progression to end-stage renal disease, in HIV-infected individuals: a tale of two races. *J Infect Dis*. 2008; 197: 1548-1557.
 95. D'Agati V, Appel GB. Renal pathology of human immunodeficiency virus infection. *Semin Nephrol*. 1998; 18: 406-421.
 96. Vardinon N, Yust I, Katz O, Iaina A, Katzir Z, Modai D, et al. Anti-HIV indeterminate western blot in dialysis patients: a long-term follow-up. *Am J Kidney Dis*. 1999; 34: 146-149.
 97. El Sayed NM, Gomatos PJ, Beck-Sague CM, Dietrich U, von Briesen H, Osmanov S, et al. Epidemic transmission of human immunodeficiency virus in renal dialysis centers in Egypt. *J Infect Dis*. 2000; 181: 91-97.
 98. Velandia M, Fridkin SK, Cardenas V, Boshell J, Ramirez G, Bland L, et al. Transmission of HIV in dialysis centre. *Lancet*. 1995; 345: 1417-1422.
 99. Henderson DK, Fahey BJ, Willy M, Schmitt JM, Carey K, Koziol DE, et al. Risk for occupational transmission of human immunodeficiency virus type 1 (HIV-1) associated with clinical exposures. A prospective evaluation. *Ann Intern Med*. 1990; 113: 740-746.
 100. Bell DM. Occupational risk of human immunodeficiency virus infection in healthcare workers: an overview. *Am J Med*. 1997; 102: 9-15.
 101. Cardo DM, Culver DH, Ciesielski CA, Srivastava PU, Marcus R, Abiteboul D, et al. A case-control study of HIV seroconversion in health care workers after percutaneous exposure. Centers for Disease Control and Prevention Needlestick Surveillance Group. *N Engl J Med*. 1997; 337: 1485-1490.
 102. Smith DK, Grohskopf LA, Black RJ, Auerbach JD, Veronese F, Struble KA, et al. Antiretroviral postexposure prophylaxis after sexual, injection-drug use, or other nonoccupational exposure to HIV in the United States: recommendations from the U.S. Department of Health and Human Services. *MMWR Recomm Rep*. 2005; 54: 1-20.
 103. Panlilio AL, Cardo DM, Grohskopf LA, Heneine W, Ross CS. Updated U.S. Public Health Service guidelines for the management of occupational exposures to HIV and recommendations for postexposure prophylaxis. *MMWR Recomm Rep*. 2005; 54: 1-17.
 104. Curi MA, Pappas PJ, Silva MB, Jr., Patel S, Padberg FT, Jr., Jamil Z, et al. Hemodialysis access: influence of the human immunodeficiency virus on patency and infection rates. *J Vasc Surg*. 1999; 29: 608-616.
 105. Mitchell D, Krishnasami Z, Young CJ, Allon M. Arteriovenous access outcomes in haemodialysis patients with HIV infection. *Nephrol Dial Transplant*. 2007; 22: 465-470.

106. Mitchell D, Krishnasami Z, Allon M. Catheter-related bacteraemia in haemodialysis patients with HIV infection. *Nephrol Dial Transplant*. 2006; 21: 3185-3188.
107. Shrivastava D, Rao TK, Sinert R, Khurana E, Lundin AP, Friedman EA. The efficacy of erythropoietin in human immunodeficiency virus-infected end-stage renal disease patients treated by maintenance hemodialysis. *Am J Kidney Dis*. 1995; 25: 904-909.
108. Oberai PC, Dalal D, Zhang L, Wang C, Eustace J, Parekh RS. Incidence of atherosclerotic cardiovascular disease among HIV patients receiving dialysis. *Am J Kidney Dis*. 2006; 47: 848-855.
109. Ahuja TS, Borucki M, Grady J. Highly active antiretroviral therapy improves survival of HIV-infected hemodialysis patients. *Am J Kidney Dis*. 2000; 36: 574-580.
110. Ortiz C, Meneses R, Jaffe D, Fernandez JA, Perez G, Bourgoignie JJ. Outcome of patients with human immunodeficiency virus on maintenance hemodialysis. *Kidney Int*. 1988; 34: 248-253.
111. Perinbasekar S, Brod-Miller C, Pal S, Mattana J. Predictors of survival in HIV-infected patients on hemodialysis. *Am J Nephrol*. 1996; 16: 280-286.
112. Ahuja TS, Grady J, Khan S. Changing trends in the survival of dialysis patients with human immunodeficiency virus in the United States. *J Am Soc Nephrol*. 2002; 13: 1889-1893.
113. Tourret J, Tostivint I, du Montcel ST, Bragg-Gresham J, Karie S, Vigneau C, et al. Outcome and prognosis factors in HIV-infected hemodialysis patients. *Clin J Am Soc Nephrol*. 2006; 1: 1241-1247.
114. Hammer SM, Eron JJ, Jr., Reiss P, Schooley RT, Thompson MA, Walmsley S, et al. Antiretroviral treatment of adult HIV infection: 2008 recommendations of the International AIDS Society-USA panel. *JAMA*. 2008; 300: 555-570.
115. Tourret J, Tostivint I, Tezenas Du Montcel S, Karie S, Launay-Vacher V, Vigneau C, et al. Antiretroviral drug dosing errors in HIV-infected patients undergoing hemodialysis. *Clin Infect Dis*. 2007; 45: 779-784.
116. Choi AI, Rodriguez RA, Bacchetti P, Volberding PA, Havlir D, Bertenthal D, et al. Low rates of antiretroviral therapy among HIV-infected patients with chronic kidney disease. *Clin Infect Dis*. 2007; 45: 1633-1639.